How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.** Do **NOT** use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months				
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL	2.5 mL		
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL [<u>***</u> *	4 mL		
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL	5 mL →	1 tablet	
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL	15stc -12.5stc 10stc -7.5stc 5stc -7.5stc	1 ½ tablets	
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL	10 mL → 15 mL −12.5 mL 10 mL	2 tablets	1 tablet
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years		-150iL-12.5ci 100iL-12.5ci -5ciL-7.5ciL-	2 ½ tablets	1 tablet
72 to 95 pounds (about 33 to 43 kilograms)	11 years		15 mL →	3 tablets	1 ½ tablets
96 pounds or more (44 kilograms or more)	12 years or older		10 mL + 10 mL = 20 mL 10 mL → 15 ot12.5 ot. and 10 mL → 15 ot12.5 ot. 10 mL → 15 ot12.5 ot. -15 ot12.5 ot15 ot12.5 ot15 ot7.5 ot.	4 tablets	2 tablets